

INTRODUCTION TO PHENOMENOLOGY WITH MASSIVE NEUTRINOS IN 2024

Concha Gonzalez-Garcia

(YITP-Stony Brook & ICREA-University of Barcelona)

60th International School of Subnuclear Physics

Erice, June 15-16th, 2024



INTRODUCTION TO PHENOMENOLOGY WITH MASSIVE NEUTRINOS IN 2024

Concha Gonzalez-Garcia

(ICREA-University of Barcelona & YITP-Stony Brook)

OUTLINE

- Historic Introduction to the SM of Massless Neutrinos
- Introducing ν mass: Dirac vs Majorana, Lepton mixing, Flavour Oscillations
- Summary of Flavour Oscillation Observations
- Status of 3ν global description
- Explorations beyond 3ν 's: steriles, NSI's, Z's...

Discovery of ν 's

- At end of 1800's radioactivity was discovered and three types identified: α , β , γ
 β : an electron comes out of the radioactive nucleus.
- Energy conservation $\Rightarrow e^-$ should have had a fixed energy


$$(A, Z) \rightarrow (A, Z + 1) + e^- \Rightarrow E_e = M(A, Z + 1) - M(A, Z)$$

Discovery of ν 's

- At end of 1800's radioactivity was discovered and three types identified: α , β , γ
 β : an electron comes out of the radioactive nucleus.

- Energy conservation $\Rightarrow e^-$ should have had a fixed energy

But 1914 **James Chadwick** showed that $(A, Z) \rightarrow (A, Z + 1) + e^- \Rightarrow E_e = M(A, Z + 1) - M(A, Z)$ the electron energy spectrum is continuous




Discovery of ν 's

- At end of 1800's radioactivity was discovered and three types identified: α , β , γ
 β : an electron comes out of the radioactive nucleus.

- Energy conservation $\Rightarrow e^-$ should have had a fixed energy

But 1914 **James Chadwick** showed that $(A, Z) \rightarrow (A, Z + 1) + e^- \Rightarrow E_e = M(A, Z + 1) - M(A, Z)$ the electron energy spectrum is continuous



Do we throw away the energy conservation?

Bohr: *we have no argument, either empirical or theoretical, for upholding the energy principle in the case of β ray disintegrations*

Discovery of ν 's

- The idea of the **neutrino** came in 1930, when **W. Pauli** tried a desperate saving operation of "the energy conservation principle".



In his letter addressed to the *Liebe Radioaktive Damen und Herren* (Dear Radioactive Ladies and Gentlemen), the participants of a meeting in Tübingen. He put forward the hypothesis that a new particle exists as *constituent of nuclei, the neutron ν* , able to explain the continuous spectrum of nuclear beta decay

$$(A, Z) \rightarrow (A, Z+1) + e^- + \nu$$

Discovery of ν 's

- The idea of the **neutrino** came in 1930, when **W. Pauli** tried a desperate saving operation of "the energy conservation principle".



In his letter addressed to the *Liebe Radioaktive Damen und Herren* (Dear Radioactive Ladies and Gentlemen), the participants of a meeting in Tübingen. He put forward the hypothesis that a new particle exists as *constituent of nuclei, the neutron ν* , able to explain the continuous spectrum of nuclear beta decay

$$(A, Z) \rightarrow (A, Z+1) + e^- + \nu$$


- The ν is **light** (in Pauli's words: m_ν should be of the same order as the m_e), **neutral** and has **spin 1/2**

Neutrino Detection

Fighting Pauli's "Curse":

I have done a terrible thing, I have postulated a particle that cannot be detected.


Sources of ν 's




The Big Bang

$$\rho_\nu = 330/\text{cm}^3$$

$$p_\nu = 0.0004 \text{ eV}$$



$E_\nu \sim \text{MeV}$



ExtraGalactic

$$E_\nu \gtrsim 30 \text{ TeV}$$

The Sun


ν_e

$$\Phi_\nu^{Earth} = 6 \times 10^{10} \nu/\text{cm}^2\text{s}$$

$$E_\nu \sim 0.1\text{--}20 \text{ MeV}$$


Human Body

$$\Phi_\nu = 340 \times 10^6 \nu/\text{day}$$



Nuclear Reactors


$$E_\nu \sim \text{few MeV}$$



Atmospheric ν 's

$$\nu_e, \nu_\mu, \bar{\nu}_e, \bar{\nu}_\mu$$

$$\Phi_\nu \sim 1 \nu/\text{cm}^2\text{s}$$



Earth's radioactivity

$$\Phi_\nu \sim 6 \times 10^6 \nu/\text{cm}^2\text{s}$$

Accelerators

$$E_\nu \simeq 0.3\text{--}30 \text{ GeV}$$



NSS

$$E_\nu \sim \text{MeV}$$

Neutrino Detection

Problem: Quantitatively: a ν sees a proton of area:

$$\sigma^{\nu p} \sim 10^{-38} \text{cm}^2 \frac{E_\nu}{\text{GeV}}$$

Neutrino Detection

Problem: Quantitatively: a ν sees a proton of area:

$$\sigma^{\nu p} \sim 10^{-38} \text{cm}^2 \frac{E_\nu}{\text{GeV}}$$

- So let's consider the atmospheric ν 's:

$$\Phi_\nu^{\text{ATM}} = 1 \nu / (\text{cm}^2 \text{ second}) \quad \text{y} \quad \langle E_\nu \rangle = 1 \text{ GeV}$$

- How many interact?

Neutrino Detection

Problem: Quantitatively: a ν sees a proton of area:

$$\sigma^{\nu p} \sim 10^{-38} \text{cm}^2 \frac{E_\nu}{\text{GeV}}$$

- So let's consider the atmospheric ν 's:

$$\Phi_\nu^{\text{ATM}} = 1 \nu / (\text{cm}^2 \text{ second}) \quad \text{y} \quad \langle E_\nu \rangle = 1 \text{ GeV}$$

- How many interact? In a human body

$$N_{\text{int}} = \Phi_\nu \times \sigma^{\nu p} \times N_{\text{prot}}^{\text{human}} \times T_{\text{life}}^{\text{human}}$$

$$\left. \begin{aligned} N_{\text{protons}}^{\text{human}} &= \frac{M^{\text{human}}}{m_p} \times N_A = 80 \text{kg} \times N_A \sim 5 \times 10^{28} \text{protons} \\ T^{\text{human}} &= 80 \text{years} = 2 \times 10^9 \text{sec} \end{aligned} \right\} \begin{aligned} & (M \times T \equiv \text{Exposure}) \\ & \text{Exposure}_{\text{human}} \\ & \sim \text{Ton} \times \text{year} \end{aligned}$$

Neutrino Detection

Problem: Quantitatively: a ν sees a proton of area:

$$\sigma^{\nu p} \sim 10^{-38} \text{cm}^2 \frac{E_\nu}{\text{GeV}}$$

- So let's consider the atmospheric ν 's:

$$\Phi_\nu^{\text{ATM}} = 1 \nu / (\text{cm}^2 \text{ second}) \quad y \quad \langle E_\nu \rangle = 1 \text{ GeV}$$

- How many interact? In a human body

$$N_{\text{int}} = \Phi_\nu \times \sigma^{\nu p} \times N_{\text{prot}}^{\text{human}} \times T_{\text{life}}^{\text{human}}$$

$$\left. \begin{aligned} N_{\text{protons}}^{\text{human}} &= \frac{M^{\text{human}}}{g_r} \times N_A = 80\text{kg} \times N_A \sim 5 \times 10^{28} \text{ protons} \\ T^{\text{human}} &= 80 \text{ years} = 2 \times 10^9 \text{ sec} \end{aligned} \right\} \begin{aligned} &(M \times T \equiv \text{Exposure}) \\ &\text{Exposure}_{\text{human}} \\ &\sim \text{Ton} \times \text{year} \end{aligned}$$

$$N_{\text{int}} = (5 \times 10^{28}) (2 \times 10^9) \times 10^{-38} \sim 1 \text{ interaction in life}$$

Neutrino Detection

Problem: Quantitatively: a ν sees a proton of area:

$$\sigma^{\nu p} \sim 10^{-38} \text{cm}^2 \frac{E_\nu}{\text{GeV}}$$

- So let's consider the atmospheric ν 's:

$$\Phi_\nu^{\text{ATM}} = 1 \nu / (\text{cm}^2 \text{ second}) \quad y \quad \langle E_\nu \rangle = 1 \text{ GeV}$$

- How many interact? In a human body

$$N_{\text{int}} = \Phi_\nu \times \sigma^{\nu p} \times N_{\text{prot}}^{\text{human}} \times T_{\text{life}}^{\text{human}}$$

$$\left. \begin{aligned} N_{\text{protons}}^{\text{human}} &= \frac{M^{\text{human}}}{m_p} \times N_A = 80 \text{kg} \times N_A \sim 5 \times 10^{28} \text{ protons} \\ T^{\text{human}} &= 80 \text{ years} = 2 \times 10^9 \text{ sec} \end{aligned} \right\} \begin{aligned} & (M \times T \equiv \text{Exposure}) \\ & \text{Exposure}_{\text{human}} \\ & \sim \text{Ton} \times \text{year} \end{aligned}$$



$$N_{\text{int}} = (5 \times 10^{28}) (2 \times 10^9) \times 10^{-38} \sim 1 \text{ interaction in life}$$

To detect neutrinos we need very intense source and/or
a huge detector with Exposure $\sim \text{Kton} \times \text{year}$

First Neutrino Detection

In 1953 **Frederick Reines** and **Clyde Cowan** put a detector near a nuclear reactor (**the most intense source available**)

400 l of water
and Cadmium Chloride.



e^+ annihilates with e^- in the water and produces **two γ 's simultaneously**.

neutron is captured by **proton** the cadmium and a γ 's is emitted **15 msec later**

Reines y Clyde saw clearly this signature: **the first neutrino had been detected**

Neutrinos = “Left-handed”


Helicity of Neutrinos*

M. GOLDHABER, L. GRODZINS, AND A. W. SUNYAR

Brookhaven National Laboratory, Upton, New York

(Received December 11, 1957)

A COMBINED analysis of circular polarization and resonant scattering of γ rays following orbital electron capture measures the helicity of the neutrino. We have carried out such a measurement with Eu^{152m} , which decays by orbital electron capture. If we assume the most plausible spin-parity assignment for this isomer compatible with its decay scheme,¹ 0^- , we find that the neutrino is “left-handed,” i.e., $\sigma_\nu \cdot \hat{p}_\nu = -1$ (negative helicity).



- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1 - \gamma_5}{2} \psi \quad \psi_R = \frac{1 + \gamma_5}{2} \psi$

- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1-\gamma_5}{2} \psi \quad \psi_R = \frac{1+\gamma_5}{2} \psi$
- The Hamiltonian for a massive fermion ψ is $H = \bar{\psi}(x) \left(-i \sum_j \gamma^j \partial_j + m \right) \psi(x)$
- 4 states with (E, \vec{p}) $(\gamma^\mu p_\mu - m) u_s(\vec{p}) = 0 \quad (\gamma^\mu p_\mu + m) v_s(\vec{p}) = 0 \quad s = 1, 2$

- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1-\gamma_5}{2} \psi \quad \psi_R = \frac{1+\gamma_5}{2} \psi$
- The Hamiltonian for a massive fermion ψ is $H = \bar{\psi}(x) \left(-i \sum_j \gamma^j \partial_j + m \right) \psi(x)$
- 4 states with (E, \vec{p}) $(\gamma^\mu p_\mu - m) u_s(\vec{p}) = 0 \quad (\gamma^\mu p_\mu + m) v_s(\vec{p}) = 0 \quad s = 1, 2$
- Since $[H, \gamma_5] \neq 0$ and $[\vec{P}, \vec{J}] \neq 0 \quad [\vec{J} = \vec{L} + \frac{\vec{\Sigma}}{2} \quad (\Sigma^i = -\gamma^0 \gamma^5 \gamma^i)]$
 \Rightarrow Neither Chirality nor J_i can characterize the fermion simultaneously with E, \vec{p}

- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1-\gamma_5}{2} \psi \quad \psi_R = \frac{1+\gamma_5}{2} \psi$
- The Hamiltonian for a massive fermion ψ is $H = \bar{\psi}(x) \left(-i \sum_j \gamma^j \partial_j + m \right) \psi(x)$
- 4 states with (E, \vec{p}) $(\gamma^\mu p_\mu - m)u_s(\vec{p}) = 0 \quad (\gamma^\mu p_\mu + m)v_s(\vec{p}) = 0 \quad s = 1, 2$
- Since $[H, \gamma_5] \neq 0$ and $[\vec{P}, \vec{J}] \neq 0 \quad [\vec{J} = \vec{L} + \frac{\vec{\Sigma}}{2} \quad (\Sigma^i = -\gamma^0 \gamma^5 \gamma^i)]$
 \Rightarrow Neither Chirality nor J_i can characterize the fermion simultaneously with E, \vec{p}
- But $[H, \vec{J} \cdot \vec{P}] = [\vec{P}, \vec{J} \cdot \vec{P}] = 0 \Rightarrow$ we can choose $u_1(\vec{p}) \equiv u_+(\vec{p})$ and $u_2(\vec{p}) \equiv u_-(\vec{p})$ (same for $v_{1,2}$) to be eigenstates of the **helicity** projector

$$\mathcal{P}_\pm = \frac{1}{2} \left(1 \pm 2 \vec{J} \frac{\vec{P}}{|\vec{P}|} \right) = \frac{1}{2} \left(1 \pm \vec{\Sigma} \frac{\vec{P}}{|\vec{P}|} \right)$$

- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1-\gamma_5}{2} \psi \quad \psi_R = \frac{1+\gamma_5}{2} \psi$
- The Hamiltonian for a massive fermion ψ is $H = \bar{\psi}(x) \left(-i \sum_j \gamma^j \partial_j + m \right) \psi(x)$
- 4 states with (E, \vec{p}) $(\gamma^\mu p_\mu - m) u_s(\vec{p}) = 0 \quad (\gamma^\mu p_\mu + m) v_s(\vec{p}) = 0 \quad s = 1, 2$
- Since $[H, \gamma_5] \neq 0$ and $[\vec{P}, \vec{J}] \neq 0 \quad [\vec{J} = \vec{L} + \frac{\vec{\Sigma}}{2} \quad (\Sigma^i = -\gamma^0 \gamma^5 \gamma^i)]$
 \Rightarrow Neither Chirality nor J_i can characterize the fermion simultaneously with E, \vec{p}
- But $[H, \vec{J} \cdot \vec{P}] = [\vec{P}, \vec{J} \cdot \vec{P}] = 0 \Rightarrow$ we can choose $u_1(\vec{p}) \equiv u_+(\vec{p})$ and $u_2(\vec{p}) \equiv u_-(\vec{p})$ (same for $v_{1,2}$) to be eigenstates of the **helicity** projector

$$\mathcal{P}_\pm = \frac{1}{2} \left(1 \pm 2 \vec{J} \frac{\vec{P}}{|\vec{P}|} \right) = \frac{1}{2} \left(1 \pm \vec{\Sigma} \frac{\vec{P}}{|\vec{P}|} \right)$$

- For **massless** fermions the Dirac equation can be written

$$\vec{\Sigma} \cdot \vec{P} \psi = -\gamma^0 \gamma^5 \vec{\gamma} \cdot \vec{p} \psi = -\gamma^0 \gamma^5 \gamma^0 E \psi = \gamma^5 E \psi \Rightarrow \text{For } m = 0 \quad \mathcal{P}_\pm = \mathcal{P}_{R,L}$$

Only for **massless** fermions **Helicity** and **chirality** states are the same.

- We define the **chiral** projections $\mathcal{P}_{R,L} = \frac{1 \pm \gamma_5}{2} \Rightarrow \psi_L = \frac{1-\gamma_5}{2} \psi \quad \psi_R = \frac{1+\gamma_5}{2} \psi$
- The Hamiltonian for a massive fermion ψ is $H = \bar{\psi}(x) \left(-i \sum_j \gamma^j \partial_j + m \right) \psi(x)$
- 4 states with (E, \vec{p}) $(\gamma^\mu p_\mu - m) u_s(\vec{p}) = 0 \quad (\gamma^\mu p_\mu + m) v_s(\vec{p}) = 0 \quad s = 1, 2$
- Since $[H, \gamma_5] \neq 0$ and $[\vec{P}, \vec{J}] \neq 0 \quad \left[\vec{J} = \vec{L} + \frac{\vec{\Sigma}}{2} \quad (\Sigma^i = -\gamma^0 \gamma^5 \gamma^i) \right]$
 \Rightarrow Neither Chirality nor J_i can characterize the fermion simultaneously with E, \vec{p}
- But $[H, \vec{J} \cdot \vec{P}] = [\vec{P}, \vec{J} \cdot \vec{P}] = 0 \Rightarrow$ we can choose $u_1(\vec{p}) \equiv u_+(\vec{p})$ and $u_2(\vec{p}) \equiv u_-(\vec{p})$ (same for $v_{1,2}$) to be eigenstates of the **helicity** projector

$$\mathcal{P}_\pm = \frac{1}{2} \left(1 \pm 2 \vec{J} \frac{\vec{P}}{|\vec{P}|} \right) = \frac{1}{2} \left(1 \pm \vec{\Sigma} \frac{\vec{P}}{|\vec{P}|} \right) = \mathcal{P}_{L,R} + \mathcal{O}\left(\frac{m}{p}\right)$$

- For **massless** fermions the Dirac equation can be written

$$\vec{\Sigma} \cdot \vec{P} \psi = -\gamma^0 \gamma^5 \vec{\gamma} \cdot \vec{p} \psi = -\gamma^0 \gamma^5 \gamma^0 E \psi = \gamma^5 E \psi \Rightarrow \text{For } m = 0 \quad \mathcal{P}_\pm = \mathcal{P}_{R,L}$$

Only for **massless** fermions **Helicity** and **chirality** states are the same.

ν in the SM

- The SM is a gauge theory based on the symmetry group

$$SU(3)_C \times SU(2)_L \times U(1)_Y \Rightarrow SU(3)_C \times U(1)_{EM}$$

- 3 Generations of Fermions:

$(1, 2, -\frac{1}{2})$	$(3, 2, \frac{1}{6})$	$(1, 1, -1)$	$(3, 1, \frac{2}{3})$	$(3, 1, -\frac{1}{3})$
L_L	Q_L^i	E_R	U_R^i	D_R^i
$\begin{pmatrix} \nu_e \\ e \\ \nu_\mu \\ \mu \\ \nu_\tau \\ \tau \end{pmatrix}_L$	$\begin{pmatrix} u^i \\ d^i \\ c^i \\ s^i \\ t^i \\ b^i \end{pmatrix}_L$	$\begin{pmatrix} e_R \\ \mu_R \\ \tau_R \end{pmatrix}$	$\begin{pmatrix} u_R^i \\ c_R^i \\ t_R^i \end{pmatrix}$	$\begin{pmatrix} d_R^i \\ s_R^i \\ b_R^i \end{pmatrix}$

- Spin-0 particle ϕ : $(1, 2, \frac{1}{2})$

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \xrightarrow{SSB} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h \end{pmatrix}$$

ν in the SM

- The SM is a gauge theory based on the symmetry group

$$SU(3)_C \times SU(2)_L \times U(1)_Y \Rightarrow SU(3)_C \times U(1)_{EM}$$

- 3 Generations of Fermions:

$(1, 2, -\frac{1}{2})$	$(3, 2, \frac{1}{6})$	$(1, 1, -1)$	$(3, 1, \frac{2}{3})$	$(3, 1, -\frac{1}{3})$
L_L	Q_L^i	E_R	U_R^i	D_R^i
$\begin{pmatrix} \nu_e \\ e \\ \nu_\mu \\ \mu \\ \nu_\tau \\ \tau \end{pmatrix}_L$	$\begin{pmatrix} u^i \\ d^i \\ c^i \\ s^i \\ t^i \\ b^i \end{pmatrix}_L$	e_R μ_R τ_R	u_R^i c_R^i t_R^i	d_R^i s_R^i b_R^i

$$Q_{EM} = T_{L3} + Y$$

- ν 's are $T_{L3} = \frac{1}{2}$ components of L_L
- ν 's have no strong or EM interactions
- No ν_R (\equiv singlets of gauge group)

- Spin-0 particle ϕ : $(1, 2, \frac{1}{2})$

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \xrightarrow{SSB} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h \end{pmatrix}$$

SM Fermion Lagrangian

$$\begin{aligned}
\mathcal{L} = & \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{Q_{L,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g \frac{\tau_a}{2} \delta_{ij} W_\mu^a - g' \frac{1}{6} \delta_{ij} B_\mu \right) Q_{L,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{U_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g' \frac{2}{3} \delta_{ij} B_\mu \right) U_{R,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{D_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a + g' \frac{1}{3} \delta_{ij} B_\mu \right) D_{R,k}^j \\
& + \sum_{k=1}^3 \overline{L_{L,k}} \gamma^\mu \left(i\partial_\mu - g \frac{\tau_i}{2} W_\mu^i + g' \frac{1}{2} B_\mu \right) L_{L,k} + \overline{E_{R,k}} \gamma^\mu (i\partial_\mu + g' B_\mu) E_{R,k} \\
& - \sum_{k,k'=1}^3 \left(\lambda_{kk'}^u \overline{Q}_{L,k} (i\tau_2) \phi^* U_{R,k'} + \lambda_{kk'}^d \overline{Q}_{L,k} \phi D_{R,k'} + \lambda_{kk'}^l \overline{L}_{L,k} \phi E_{R,k'} + h.c. \right)
\end{aligned}$$

SM Fermion Lagrangian

$$\begin{aligned}
\mathcal{L} = & \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{Q_{L,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g \frac{\tau_a}{2} \delta_{ij} W_\mu^a - g' \frac{1}{6} \delta_{ij} B_\mu \right) Q_{L,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{U_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g' \frac{2}{3} \delta_{ij} B_\mu \right) U_{R,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{D_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a + g' \frac{1}{3} \delta_{ij} B_\mu \right) D_{R,k}^j \\
& + \sum_{k=1}^3 \overline{L_{L,k}} \gamma^\mu \left(i\partial_\mu - g \frac{\tau_i}{2} W_\mu^i + g' \frac{1}{2} B_\mu \right) L_{L,k} + \overline{E_{R,k}} \gamma^\mu (i\partial_\mu + g' B_\mu) E_{R,k} \\
& - \sum_{k,k'=1}^3 \left(\lambda_{kk'}^u \overline{Q}_{L,k} (i\tau_2) \phi^* U_{R,k'} + \lambda_{kk'}^d \overline{Q}_{L,k} \phi D_{R,k'} + \lambda_{kk'}^l \overline{L}_{L,k} \phi E_{R,k'} + h.c. \right)
\end{aligned}$$

- Invariant under global rotations

$$Q_{L,k} \rightarrow e^{i\alpha_B/3} Q_{L,k} \quad U_{R,k} \rightarrow e^{i\alpha_B/3} U_{R,k} \quad D_{R,k} \rightarrow e^{i\alpha_B/3} D_{R,k} \quad L_{L,k} \rightarrow e^{i\alpha_{L_k}} L_{L,k} \quad E_{R,k} \rightarrow e^{i\alpha_{L_k}} E_{R,k}$$

SM Fermion Lagrangian

$$\begin{aligned}
\mathcal{L} = & \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{Q_{L,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g \frac{\tau_a}{2} \delta_{ij} W_\mu^a - g' \frac{1}{6} \delta_{ij} B_\mu \right) Q_{L,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{U_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a - g' \frac{2}{3} \delta_{ij} B_\mu \right) U_{R,k}^j \\
& + \sum_{k=1}^3 \sum_{i,j=1}^3 \overline{D_{R,k}^i} \gamma^\mu \left(i\partial_\mu - g_s \frac{\lambda_{a,ij}}{2} G_\mu^a + g' \frac{1}{3} \delta_{ij} B_\mu \right) D_{R,k}^j \\
& + \sum_{k=1}^3 \overline{L_{L,k}} \gamma^\mu \left(i\partial_\mu - g \frac{\tau_i}{2} W_\mu^i + g' \frac{1}{2} B_\mu \right) L_{L,k} + \overline{E_{R,k}} \gamma^\mu (i\partial_\mu + g' B_\mu) E_{R,k} \\
& - \sum_{k,k'=1}^3 \left(\lambda_{kk'}^u \overline{Q}_{L,k} (i\tau_2) \phi^* U_{R,k'} + \lambda_{kk'}^d \overline{Q}_{L,k} \phi D_{R,k'} + \lambda_{kk'}^l \overline{L}_{L,k} \phi E_{R,k'} + h.c. \right)
\end{aligned}$$

- Invariant under global rotations

$$Q_{L,k} \rightarrow e^{i\alpha_B/3} Q_{L,k} \quad U_{R,k} \rightarrow e^{i\alpha_B/3} U_{R,k} \quad D_{R,k} \rightarrow e^{i\alpha_B/3} D_{R,k} \quad L_{L,k} \rightarrow e^{i\alpha_{L_k}} L_{L,k} \quad E_{R,k} \rightarrow e^{i\alpha_{L_k}} E_{R,k}$$

\Rightarrow **Accidental** (\equiv *not imposed*) global **symmetry**: $U(1)_B \times U(1)_{L_e} \times U(1)_{L_\mu} \times U(1)_{L_\tau}$

\Rightarrow **Each lepton flavour**, L_i , is conserved

\Rightarrow **Total lepton number** $L = L_e + L_\mu + L_\tau$ is conserved

- A **fermion mass** can be seen as at a **Left-Right** transition

$$m_f \bar{\psi} \psi = m_f \bar{\psi}_L \psi_R + h.c. \quad (\text{this is not } SU(2)_L \text{ gauge invariant})$$

- A **fermion mass** can be seen as at a **Left-Right transition**

$$m_f \bar{\psi} \psi = m_f \bar{\psi}_L \psi_R + h.c. \quad (\text{this is not } SU(2)_L \text{ gauge invariant})$$

- In the Standard Model mass comes from *spontaneous symmetry breaking* via Yukawa interaction of the left-handed doublet L_L with the right-handed singlet E_R :

$$\mathcal{L}_Y^l = -\lambda_{ij}^l \bar{L}_{Li} E_{Rj} \phi + h.c. \quad \phi = \text{the scalar doublet}$$

- A **fermion mass** can be seen as at a **Left-Right transition**

$$m_f \bar{\psi} \psi = m_f \bar{\psi}_L \psi_R + h.c. \quad (\text{this is not } SU(2)_L \text{ gauge invariant})$$

- In the Standard Model mass comes from *spontaneous symmetry breaking* via Yukawa interaction of the left-handed doublet L_L with the right-handed singlet E_R :

$$\mathcal{L}_Y^l = -\lambda_{ij}^l \bar{L}_{Li} E_{Rj} \phi + h.c. \quad \phi = \text{the scalar doublet}$$

- After spontaneous symmetry breaking

$$\phi \xrightarrow{SSB} \begin{pmatrix} 0 \\ \frac{v+h}{\sqrt{2}} \end{pmatrix} \Rightarrow \mathcal{L}_{\text{mass}}^l = -\bar{E}_L M^\ell E_R + h.c. \text{ with } M^\ell = \frac{1}{\sqrt{2}} \lambda^l v$$

- A **fermion mass** can be seen as at a **Left-Right transition**

$$m_f \bar{\psi} \psi = m_f \bar{\psi}_L \psi_R + h.c. \quad (\text{this is not } SU(2)_L \text{ gauge invariant})$$

- In the Standard Model mass comes from *spontaneous symmetry breaking* via Yukawa interaction of the left-handed doublet L_L with the right-handed singlet E_R :

$$\mathcal{L}_Y^l = -\lambda_{ij}^l \bar{L}_{Li} E_{Rj} \phi + h.c. \quad \phi = \text{the scalar doublet}$$

- After spontaneous symmetry breaking

$$\phi \xrightarrow{SSB} \begin{pmatrix} 0 \\ \frac{v+h}{\sqrt{2}} \end{pmatrix} \Rightarrow \mathcal{L}_{\text{mass}}^l = -\bar{E}_L M^\ell E_R + h.c. \text{ with } M^\ell = \frac{1}{\sqrt{2}} \lambda^l v$$

In the SM:

- There are no right-handed neutrinos
 \Rightarrow No renormalizable (ie $\dim \leq 4$) gauge-invariant operator for tree level ν mass
- SM gauge invariance \Rightarrow accidental symmetry $U(1)_B \times U(1)_{L_e} \times U(1)_{L_\mu} \times U(1)_{L_\tau}$
 \Rightarrow Not possible to generate such term at any order perturbatively

- A **fermion mass** can be seen as at a **Left-Right transition**

$$m_f \bar{\psi} \psi = m_f \bar{\psi}_L \psi_R + h.c. \quad (\text{this is not } SU(2)_L \text{ gauge invariant})$$

- In the Standard Model mass comes from *spontaneous symmetry breaking* via Yukawa interaction of the left-handed doublet L_L with the right-handed singlet E_R :

$$\mathcal{L}_Y^l = -\lambda_{ij}^l \bar{L}_{Li} E_{Rj} \phi + h.c. \quad \phi = \text{the scalar doublet}$$

- After spontaneous symmetry breaking

$$\phi \xrightarrow{SSB} \begin{pmatrix} 0 \\ \frac{v+h}{\sqrt{2}} \end{pmatrix} \Rightarrow \mathcal{L}_{\text{mass}}^l = -\bar{E}_L M^\ell E_R + h.c. \text{ with } M^\ell = \frac{1}{\sqrt{2}} \lambda^l v$$

In the SM:

- There are no right-handed neutrinos
 \Rightarrow No renormalizable (ie $\dim \leq 4$) gauge-invariant operator for tree level ν mass
- SM gauge invariance \Rightarrow accidental symmetry $U(1)_B \times U(1)_{L_e} \times U(1)_{L_\mu} \times U(1)_{L_\tau}$
 \Rightarrow Not possible to generate such term at any order perturbatively

In SM ν 's are *Strictly* Massless & Lepton Flavours are *Strictly* Conserved

- We have observed with high (or good) precision:
 - * Atmospheric ν_μ & $\bar{\nu}_\mu$ disappear most likely to ν_τ (**SK**, MINOS, ICECUBE)
 - * Accel. ν_μ & $\bar{\nu}_\mu$ disappear at $L \sim 300/800$ Km (**K2K**, **T2K**, **MINOS**, **NO ν A**)
 - * Some accelerator ν_μ appear as ν_e at $L \sim 300/800$ Km (**T2K**, MINOS, NO ν A)
 - * Solar ν_e convert to ν_μ/ν_τ (**Cl**, **Ga**, **SK**, **SNO**, **Borexino**)
 - * Reactor $\bar{\nu}_e$ disappear at $L \sim 200$ Km (**KamLAND**)
 - * Reactor $\bar{\nu}_e$ disappear at $L \sim 1$ Km (**D-Chooz**, **Daya Bay**, **Reno**)

- We have observed with high (or good) precision:
 - * Atmospheric ν_μ & $\bar{\nu}_\mu$ disappear most likely to ν_τ (**SK**, MINOS, ICECUBE)
 - * Accel. ν_μ & $\bar{\nu}_\mu$ disappear at $L \sim 300/800$ Km (**K2K**, **T2K**, **MINOS**, **NO ν A**)
 - * Some accelerator ν_μ appear as ν_e at $L \sim 300/800$ Km (**T2K**, MINOS, NO ν A)
 - * Solar ν_e convert to ν_μ/ν_τ (**Cl**, **Ga**, **SK**, **SNO**, **Borexino**)
 - * Reactor $\bar{\nu}_e$ disappear at $L \sim 200$ Km (**KamLAND**)
 - * Reactor $\bar{\nu}_e$ disappear at $L \sim 1$ Km (**D-Chooz**, **Daya Bay**, **Reno**)

All this implies that L_α are violated

and There is Physics Beyond SM

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:
 - Their own antiparticle such as γ , π^0 ...
 - Different from their antiparticle such as K^0 , \bar{K}^0 ...
- In the SM ν are the only *neutral fermions*

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:
 - Their own antiparticle such as γ , π^0 ...
 - Different from their antiparticle such as K^0 , \bar{K}^0 ...
 - In the SM ν are the only *neutral fermions*
- ⇒ **OPEN QUESTION:** are neutrino and antineutrino the same or different particles?

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:

- Their own antiparticle such as γ , π^0 ...
- Different from their antiparticle such as K^0 , \bar{K}^0 ...

- In the SM ν are the only *neutral fermions*

⇒ **OPEN QUESTION:** are neutrino and antineutrino the same or different particles?

* ANSWER 1: ν different from anti- ν ⇒ ν is a *Dirac* fermion (like e)

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:
 - Their own antiparticle such as γ , π^0 ...
 - Different from their antiparticle such as K^0 , \bar{K}^0 ...

- In the SM ν are the only *neutral fermions*

⇒ **OPEN QUESTION:** are neutrino and antineutrino the same or different particles?

* **ANSWER 1:** ν different from anti- ν ⇒ ν is a *Dirac* fermion (like e)

⇒ It is described by a *Dirac* field $\nu(x) = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:
 - Their own antiparticle such as $\gamma, \pi^0 \dots$
 - Different from their antiparticle such as $K^0, \bar{K}^0 \dots$

- In the SM ν are the only *neutral fermions*

⇒ **OPEN QUESTION:** are neutrino and antineutrino the same or different particles?

* **ANSWER 1:** ν different from anti- ν ⇒ ν is a *Dirac* fermion (like e)

⇒ It is described by a *Dirac* field $\nu(x) = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$

⇒ And the charged conjugate neutrino field \equiv the antineutrino field

$$\nu^C = C \nu C^{-1} = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = -C \bar{\nu}^T$$

($C = i\gamma^2\gamma^0$)

which contain two sets of creation–annihilation operators

Dirac versus Majorana Neutrinos

- In the SM neutral bosons can be of two type:
 - Their own antiparticle such as $\gamma, \pi^0 \dots$
 - Different from their antiparticle such as $K^0, \bar{K}^0 \dots$

- In the SM ν are the only *neutral fermions*

⇒ **OPEN QUESTION:** are neutrino and antineutrino the same or different particles?

* **ANSWER 1:** ν different from anti- ν ⇒ ν is a *Dirac* fermion (like e)

⇒ It is described by a *Dirac* field $\nu(x) = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$

⇒ And the charged conjugate neutrino field \equiv the antineutrino field

$$\nu^C = C \nu C^{-1} = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = -C \bar{\nu}^T$$

$(C = i\gamma^2 \gamma^0)$

which contain two sets of creation–annihilation operators

⇒ 4 chiral fields

$$\nu_L, \nu_R, (\nu_L)^C, (\nu_R)^C \quad \text{with} \quad \nu = \nu_L + \nu_R \quad \text{and} \quad \nu^C = (\nu_L)^C + (\nu_R)^C$$

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

$$\Rightarrow \text{So we can rewrite the field } \nu_M = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

which contains only one set of creation–annihilation operators

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

$$\Rightarrow \text{So we can rewrite the field } \nu_M = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

which contains only one set of creation–annihilation operators

\Rightarrow A Majorana particle can be described with only 2 independent chiral fields:

$$\nu_L \text{ and } (\nu_L)^C \quad \text{and the other two are} \quad \nu_R = (\nu_L)^C \quad (\nu_R)^C = \nu_L$$

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

$$\Rightarrow \text{So we can rewrite the field } \nu_M = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

which contains only one set of creation–annihilation operators

\Rightarrow A Majorana particle can be described with only 2 independent chiral fields:

$$\nu_L \text{ and } (\nu_L)^C \quad \text{and the other two are} \quad \nu_R = (\nu_L)^C \quad (\nu_R)^C = \nu_L$$

• In the SM the interaction term for neutrinos

$$\mathcal{L}_{int} = \frac{i g}{\sqrt{2}} [(\bar{l}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) W_\mu^- + (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L l_\alpha) W_\mu^+] + \frac{i g}{\sqrt{2} \cos \theta_W} (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) Z_\mu$$

Only involves two chiral fields $\mathcal{P}_L \nu = \nu_L$ and $\bar{\nu} \mathcal{P}_R = (\nu_L)^{C T} C^\dagger$

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a Majorana fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

$$\Rightarrow \text{So we can rewrite the field } \nu_M = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

which contains only one set of creation-annihilation operators

\Rightarrow A Majorana particle can be described with only 2 independent chiral fields:

$$\nu_L \text{ and } (\nu_L)^C \quad \text{and the other two are} \quad \nu_R = (\nu_L)^C \quad (\nu_R)^C = \nu_L$$

• In the SM the interaction term for neutrinos

$$\mathcal{L}_{int} = \frac{i g}{\sqrt{2}} [(\bar{l}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) W_\mu^- + (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L l_\alpha) W_\mu^+] + \frac{i g}{\sqrt{2} \cos \theta_W} (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) Z_\mu$$

Only involves two chiral fields $\mathcal{P}_L \nu = \nu_L$ and $\bar{\nu} \mathcal{P}_R = (\nu_L)^{C T} C^\dagger$

\Rightarrow Weak interaction cannot distinguish if neutrinos are Dirac or Majorana

Dirac versus Majorana Neutrinos

* ANSWER 2: ν same as anti- ν $\Rightarrow \nu$ is a *Majorana* fermion : $\nu_M = \nu_M^C$

$$\Rightarrow \nu^C = \sum_{s, \vec{p}} \left[b_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right] = \nu = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + b_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

$$\Rightarrow \text{So we can rewrite the field } \nu_M = \sum_{s, \vec{p}} \left[a_s(\vec{p}) u_s(\vec{p}) e^{-i p x} + a_s^\dagger(\vec{p}) v_s(\vec{p}) e^{i p x} \right]$$

which contains only one set of creation-annihilation operators

\Rightarrow A Majorana particle can be described with only 2 independent chiral fields:

$$\nu_L \text{ and } (\nu_L)^C \quad \text{and the other two are} \quad \nu_R = (\nu_L)^C \quad (\nu_R)^C = \nu_L$$

• In the SM the interaction term for neutrinos

$$\mathcal{L}_{int} = \frac{i g}{\sqrt{2}} [(\bar{l}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) W_\mu^- + (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L l_\alpha) W_\mu^+] + \frac{i g}{\sqrt{2} \cos \theta_W} (\bar{\nu}_\alpha \gamma_\mu \mathcal{P}_L \nu_\alpha) Z_\mu$$

Only involves two chiral fields $\mathcal{P}_L \nu = \nu_L$ and $\bar{\nu} \mathcal{P}_R = (\nu_L)^{C T} C^\dagger$

\Rightarrow Weak interaction cannot distinguish if neutrinos are *Dirac or Majorana*

The difference arises when including *a neutrino mass*

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$
- One introduces ν_R which can couple to the lepton doublet by Yukawa interaction

$$\mathcal{L}_Y^{(\nu)} = -\lambda_{ij}^\nu \overline{\nu}_{Ri} L_{Lj} \tilde{\phi}^\dagger + \text{h.c.} \quad (\tilde{\phi} = i\tau_2 \phi^*)$$

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$
- One introduces ν_R which can couple to the lepton doublet by Yukawa interaction

$$\mathcal{L}_Y^{(\nu)} = -\lambda_{ij}^\nu \overline{\nu_{Ri}} L_{Lj} \tilde{\phi}^\dagger + h.c. \quad (\tilde{\phi} = i\tau_2 \phi^*)$$

- Under spontaneous symmetry-breaking $\mathcal{L}_Y^{(\nu)} \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})}$

$$\mathcal{L}_{\text{mass}}^{(\text{Dirac})} = -\overline{\nu_R} M_D^\nu \nu_L + h.c. \equiv -\frac{1}{2} (\overline{\nu_R} M_D^\nu \nu_L + \overline{(\nu_L)^c} M_D^{\nu T} (\nu_R)^c) + h.c. \equiv -\sum_k m_k \overline{\nu}_k^D \nu_k^D$$

$$M_D^\nu = \frac{1}{\sqrt{2}} \lambda^\nu v = \text{Dirac mass for neutrinos}$$

$$V_R^{\nu\dagger} M_D V^\nu = \text{diag}(m_1, m_2, m_3)$$

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$
- One introduces ν_R which can couple to the lepton doublet by Yukawa interaction

$$\mathcal{L}_Y^{(\nu)} = -\lambda_{ij}^\nu \overline{\nu_{Ri}} L_{Lj} \tilde{\phi}^\dagger + h.c. \quad (\tilde{\phi} = i\tau_2 \phi^*)$$

- Under spontaneous symmetry-breaking $\mathcal{L}_Y^{(\nu)} \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})}$

$$\mathcal{L}_{\text{mass}}^{(\text{Dirac})} = -\overline{\nu_R} M_D^\nu \nu_L + h.c. \equiv -\frac{1}{2} (\overline{\nu_R} M_D^\nu \nu_L + \overline{(\nu_L)^c} M_D^{\nu T} (\nu_R)^c) + h.c. \equiv -\sum_k m_k \overline{\nu}_k^D \nu_k^D$$

$$M_D^\nu = \frac{1}{\sqrt{2}} \lambda^\nu v = \text{Dirac mass for neutrinos}$$

$$V_R^{\nu\dagger} M_D V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_D^ν are Dirac fermions (same as quarks and charged leptons)

$$\nu^D = V^{\nu\dagger} \nu_L + V_R^{\nu\dagger} \nu_R$$

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$
- One introduces ν_R which can couple to the lepton doublet by Yukawa interaction

$$\mathcal{L}_Y^{(\nu)} = -\lambda_{ij}^\nu \overline{\nu_{Ri}} L_{Lj} \tilde{\phi}^\dagger + h.c. \quad (\tilde{\phi} = i\tau_2 \phi^*)$$

- Under spontaneous symmetry-breaking $\mathcal{L}_Y^{(\nu)} \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})}$

$$\mathcal{L}_{\text{mass}}^{(\text{Dirac})} = -\overline{\nu_R} M_D^\nu \nu_L + h.c. \equiv -\frac{1}{2} (\overline{\nu_R} M_D^\nu \nu_L + \overline{(\nu_L)^c} M_D^{\nu T} (\nu_R)^c) + h.c. \equiv -\sum_k m_k \overline{\nu}_k^D \nu_k^D$$

$$M_D^\nu = \frac{1}{\sqrt{2}} \lambda^\nu v = \text{Dirac mass for neutrinos}$$

$$V_R^{\nu\dagger} M_D V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_D^ν are Dirac fermions (same as quarks and charged leptons)

$$\nu^D = V^{\nu\dagger} \nu_L + V_R^{\nu\dagger} \nu_R$$

- $\mathcal{L}_{\text{mass}}^{(\text{Dirac})}$ involves the four chiral fields ν_L , ν_R , $(\nu_L)^c$, $(\nu_R)^c$

Adding ν Mass: Dirac Mass

- A **fermion mass** is a **Left-Right** operator : $\mathcal{L}_{m_f} = -m_f \overline{\psi}_L \psi_R + h.c.$
- One introduces ν_R which can couple to the lepton doublet by Yukawa interaction

$$\mathcal{L}_Y^{(\nu)} = -\lambda_{ij}^\nu \overline{\nu_{Ri}} L_{Lj} \tilde{\phi}^\dagger + h.c. \quad (\tilde{\phi} = i\tau_2 \phi^*)$$

- Under spontaneous symmetry-breaking $\mathcal{L}_Y^{(\nu)} \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})}$

$$\mathcal{L}_{\text{mass}}^{(\text{Dirac})} = -\overline{\nu_R} M_D^\nu \nu_L + h.c. \equiv -\frac{1}{2} (\overline{\nu_R} M_D^\nu \nu_L + \overline{(\nu_L)^c} M_D^{\nu T} (\nu_R)^c) + h.c. \equiv -\sum_k m_k \overline{\nu}_k^D \nu_k^D$$

$$M_D^\nu = \frac{1}{\sqrt{2}} \lambda^\nu v = \text{Dirac mass for neutrinos}$$

$$V_R^{\nu\dagger} M_D V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_D^ν are Dirac fermions (same as quarks and charged leptons)

$$\nu^D = V^{\nu\dagger} \nu_L + V_R^{\nu\dagger} \nu_R$$

- $\mathcal{L}_{\text{mass}}^{(\text{Dirac})}$ involves the four chiral fields ν_L , ν_R , $(\nu_L)^c$, $(\nu_R)^c$

\Rightarrow **Total Lepton number** is **conserved** by construction (not accidentally):

$$\left. \begin{array}{l} U(1)_L : \nu \rightarrow e^{i\alpha} \nu \quad \text{and} \quad \overline{\nu} \rightarrow e^{-i\alpha} \overline{\nu} \\ U(1)_L : \nu^c \rightarrow e^{-i\alpha} \nu^c \quad \text{and} \quad \overline{\nu^c} \rightarrow e^{i\alpha} \overline{\nu^c} \end{array} \right\} \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})} \rightarrow \mathcal{L}_{\text{mass}}^{(\text{Dirac})}$$

- One **does not** introduce ν_R but uses that the field $(\nu_L)^c$ is right-handed, so that one can write a **Lorentz-invariant** mass term

$$\mathcal{L}_{\text{mass}}^{(\text{Maj})} = -\frac{1}{2} \overline{\nu_L^c} M_M^\nu \nu_L + \text{h.c.} \equiv -\frac{1}{2} \sum_k m_k \bar{\nu}_i^M \nu_i^M$$

- One **does not** introduce ν_R but uses that the field $(\nu_L)^c$ is right-handed, so that one can write a **Lorentz-invariant** mass term

$$\mathcal{L}_{\text{mass}}^{(\text{Maj})} = -\frac{1}{2} \overline{\nu_L^c} M_M^\nu \nu_L + \text{h.c.} \equiv -\frac{1}{2} \sum_k m_k \bar{\nu}_i^M \nu_i^M$$

M_M^ν = Majorana mass for ν 's is *symmetric*

$$V^{\nu T} M_M V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_M^ν are Majorana particles

$$\nu^M = V^{\nu\dagger} \nu_L + (V^{\nu\dagger} \nu_L)^c \quad (\text{verify } \nu_i^{M^c} = \nu_i^M)$$

- One **does not** introduce ν_R but uses that the field $(\nu_L)^c$ is right-handed, so that one can write a **Lorentz-invariant** mass term

$$\mathcal{L}_{\text{mass}}^{(\text{Maj})} = -\frac{1}{2} \overline{\nu_L^c} M_M^\nu \nu_L + \text{h.c.} \equiv -\frac{1}{2} \sum_k m_k \bar{\nu}_i^M \nu_i^M$$

M_M^ν = Majorana mass for ν 's is *symmetric*

$$V^{\nu T} M_M V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_M^ν are Majorana particles

$$\nu^M = V^{\nu\dagger} \nu_L + (V^{\nu\dagger} \nu_L)^c \quad (\text{verify } \nu_i^{M^c} = \nu_i^M)$$

\Rightarrow **But $SU(2)_L$ gauge inv is broken** $\Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})}$ not possible at tree-level in the SM

- One **does not** introduce ν_R but uses that the field $(\nu_L)^c$ is right-handed, so that one can write a **Lorentz-invariant** mass term

$$\mathcal{L}_{\text{mass}}^{(\text{Maj})} = -\frac{1}{2} \overline{\nu_L^c} M_M^\nu \nu_L + \text{h.c.} \equiv -\frac{1}{2} \sum_k m_k \bar{\nu}_i^M \nu_i^M$$

M_M^ν = Majorana mass for ν 's is symmetric

$$V^{\nu T} M_M V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_M^ν are Majorana particles

$$\nu^M = V^{\nu\dagger} \nu_L + (V^{\nu\dagger} \nu_L)^c \quad (\text{verify } \nu_i^{Mc} = \nu_i^M)$$

\Rightarrow But $SU(2)_L$ gauge inv is broken $\Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})}$ not possible at tree-level in the SM

- Moreover under any $U(1)$ symmetry with $U(1) : \nu \rightarrow e^{i\alpha} \nu$

$$\Rightarrow \nu^c \rightarrow e^{-i\alpha} \nu^c \quad \text{and} \quad \bar{\nu} \rightarrow e^{-i\alpha} \bar{\nu} \quad \text{so} \quad \bar{\nu}^c \rightarrow e^{i\alpha} \bar{\nu}^c \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})} \rightarrow e^{2i\alpha} \mathcal{L}_{\text{mass}}^{(\text{Maj})}$$

$\mathcal{L}_{\text{mass}}^{(\text{Maj})}$ breaks $U(1) \Rightarrow$ only possible for particles without electric charge

- One **does not** introduce ν_R but uses that the field $(\nu_L)^c$ is right-handed, so that one can write a **Lorentz-invariant** mass term

$$\mathcal{L}_{\text{mass}}^{(\text{Maj})} = -\frac{1}{2} \overline{\nu_L^c} M_M^\nu \nu_L + \text{h.c.} \equiv -\frac{1}{2} \sum_k m_k \bar{\nu}_i^M \nu_i^M$$

M_M^ν = Majorana mass for ν 's is *symmetric*

$$V^{\nu T} M_M V^\nu = \text{diag}(m_1, m_2, m_3)$$

\Rightarrow The eigenstates of M_M^ν are Majorana particles

$$\nu^M = V^{\nu\dagger} \nu_L + (V^{\nu\dagger} \nu_L)^c \quad (\text{verify } \nu_i^{Mc} = \nu_i^M)$$

\Rightarrow **But $SU(2)_L$ gauge inv is broken** $\Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})}$ not possible at tree-level in the SM

- Moreover under any $U(1)$ symmetry with $U(1) : \nu \rightarrow e^{i\alpha} \nu$

$$\Rightarrow \nu^c \rightarrow e^{-i\alpha} \nu^c \quad \text{and} \quad \bar{\nu} \rightarrow e^{-i\alpha} \bar{\nu} \quad \text{so} \quad \bar{\nu}^c \rightarrow e^{i\alpha} \bar{\nu}^c \Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})} \rightarrow e^{2i\alpha} \mathcal{L}_{\text{mass}}^{(\text{Maj})}$$

$\mathcal{L}_{\text{mass}}^{(\text{Maj})}$ breaks $U(1) \Rightarrow$ only possible for particles without electric charge

\Rightarrow **Breaks Total Lepton Number** $\Rightarrow \mathcal{L}_{\text{mass}}^{(\text{Maj})}$ **not generated at any order in the SM**

ν Mass \Rightarrow Lepton Mixing

- CC and mass for 3 charged leptons ℓ_i and N neutrinos in weak basis $\nu^W \equiv \begin{pmatrix} \nu_{L,e} \\ \nu_{L,\mu} \\ \nu_{L,\tau} \\ (\nu_{R,1})^C \\ \vdots \\ \vdots \end{pmatrix}$
- $$\mathcal{L}_{CC} + \mathcal{L}_M = -\frac{g}{\sqrt{2}} \sum_{i=1}^3 \overline{\ell_{L,i}^W} \gamma^\mu \nu_i^W W_\mu^+ - \sum_{i,j=1}^3 \overline{\ell_{L,i}^W} M_{\ell ij} \ell_{R,j}^W - \frac{1}{2} \sum_{i,j=1}^N \overline{\nu_i^{cW}} M_{\nu ij} \nu_j^W + \text{h.c.}$$

ν Mass \Rightarrow Lepton Mixing

- CC and mass for 3 charged leptons ℓ_i and N neutrinos in weak basis $\nu^W \equiv \begin{pmatrix} \nu_{L,e} \\ \nu_{L,\mu} \\ \nu_{L,\tau} \\ (\nu_{R,1})^C \\ \vdots \\ \vdots \end{pmatrix}$
- $$\mathcal{L}_{CC} + \mathcal{L}_M = -\frac{g}{\sqrt{2}} \sum_{i=1}^3 \overline{\ell_{L,i}^W} \gamma^\mu \nu_i^W W_\mu^+ - \sum_{i,j=1}^3 \overline{\ell_{L,i}^W} M_{\ell ij} \ell_{R,j}^W - \frac{1}{2} \sum_{i,j=1}^N \overline{\nu_i^{cW}} M_{\nu ij} \nu_j^W + \text{h.c.}$$

- Change to mass basis : $\ell_{L,i}^W = V_{Lij}^\ell \ell_{L,j}$ $\ell_{R,i}^W = V_{Rij}^\ell \ell_{R,j}$ $\nu_i^W = V_{ij}^\nu \nu_j$

$$V_L^{\ell\dagger} M_\ell V_R^\ell = \text{diag}(m_e, m_\mu, m_\tau)$$

$$V_{L,R}^\ell \equiv \text{Unitary } 3 \times 3 \text{ matrices}$$

$$V^{\nu T} M_\nu V^\nu = \text{diag}(m_1^2, m_2^2, m_3^2, \dots, m_N^2)$$

$$V^\nu \equiv \text{Unitary } N \times N \text{ matrix.}$$

ν Mass \Rightarrow Lepton Mixing

- CC and mass for 3 charged leptons ℓ_i and N neutrinos in weak basis $\nu^W \equiv \begin{pmatrix} \nu_{L,e} \\ \nu_{L,\mu} \\ \nu_{L,\tau} \\ (\nu_{R,1})^C \\ \vdots \\ \vdots \end{pmatrix}$
- $$\mathcal{L}_{CC} + \mathcal{L}_M = -\frac{g}{\sqrt{2}} \sum_{i=1}^3 \overline{\ell_{L,i}^W} \gamma^\mu \nu_i^W W_\mu^+ - \sum_{i,j=1}^3 \overline{\ell_{L,i}^W} M_{\ell ij} \ell_{R,j}^W - \frac{1}{2} \sum_{i,j=1}^N \overline{\nu_i^{cW}} M_{\nu ij} \nu_j^W + \text{h.c.}$$

- Change to mass basis : $\ell_{L,i}^W = V_{Lij}^\ell \ell_{L,j}$ $\ell_{R,i}^W = V_{Rij}^\ell \ell_{R,j}$ $\nu_i^W = V_{ij}^\nu \nu_j$

$$V_L^{\ell\dagger} M_\ell V_R^\ell = \text{diag}(m_e, m_\mu, m_\tau)$$

$$V^{\nu T} M_\nu V^\nu = \text{diag}(m_1^2, m_2^2, m_3^2, \dots, m_N^2)$$

$V_{L,R}^\ell \equiv$ Unitary 3×3 matrices

$V^\nu \equiv$ Unitary $N \times N$ matrix.

- The charged current in the mass basis: $\mathcal{L}_{CC} = -\frac{g}{\sqrt{2}} \overline{\ell_L^i} \gamma^\mu U_{\text{LEP}}^{ij} \nu_j W_\mu^+$

ν Mass \Rightarrow Lepton Mixing

- CC and mass for 3 charged leptons ℓ_i and N neutrinos in weak basis $\nu^W \equiv \begin{pmatrix} \nu_{L,e} \\ \nu_{L,\mu} \\ \nu_{L,\tau} \\ (\nu_{R,1})^C \\ \vdots \\ \vdots \end{pmatrix}$
- $$\mathcal{L}_{CC} + \mathcal{L}_M = -\frac{g}{\sqrt{2}} \sum_{i=1}^3 \overline{\ell_{L,i}^W} \gamma^\mu \nu_i^W W_\mu^+ - \sum_{i,j=1}^3 \overline{\ell_{L,i}^W} M_{\ell ij} \ell_{R,j}^W - \frac{1}{2} \sum_{i,j=1}^N \overline{\nu_i^{cW}} M_{\nu ij} \nu_j^W + \text{h.c.}$$

- Change to mass basis : $\ell_{L,i}^W = V_{Lij}^\ell \ell_{L,j}$ $\ell_{R,i}^W = V_{Rij}^\ell \ell_{R,j}$ $\nu_i^W = V_{ij}^\nu \nu_j$

$$V_L^{\ell\dagger} M_\ell V_R^\ell = \text{diag}(m_e, m_\mu, m_\tau)$$

$$V^{\nu T} M_\nu V^\nu = \text{diag}(m_1^2, m_2^2, m_3^2, \dots, m_N^2)$$

$V_{L,R}^\ell \equiv$ Unitary 3×3 matrices

$V^\nu \equiv$ Unitary $N \times N$ matrix.

- The charged current in the mass basis: $\mathcal{L}_{CC} = -\frac{g}{\sqrt{2}} \overline{\ell_L^i} \gamma^\mu U_{\text{LEP}}^{ij} \nu_j W_\mu^+$
- $U_{\text{LEP}} \equiv 3 \times N$ matrix $U_{\text{LEP}} U_{\text{LEP}}^\dagger = I_{3 \times 3}$ but in general $U_{\text{LEP}}^\dagger U_{\text{LEP}} \neq I_{N \times N}$

$$U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^\ell V_L^{\ell\dagger ik} V^{\nu kj} P_{jj}^\nu$$

Lepton Mixing

$U_{\text{LEP}} \equiv 3 \times N$ matrix

$$U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^{\ell} V_L^{\ell \dagger ik} V^{\nu kj} P_{jj}^{\nu}$$

Lepton Mixing

$$U_{\text{LEP}} \equiv 3 \times N \text{ matrix} \quad U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^{\ell} V_L^{\ell \dagger ik} V^{\nu kj} P_{jj}^{\nu}$$

- $P_{ii}^{\ell} \supset 3$ phases absorbed in l_i
- $P_{kk}^{\nu} \supset N-1$ phases absorbed in ν_i (only possible if ν_i is Dirac)

Lepton Mixing

$$U_{\text{LEP}} \equiv 3 \times N \text{ matrix} \quad U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^{\ell} V_L^{\ell \dagger ik} V^{\nu kj} P_{jj}^{\nu}$$

- $P_{ii}^{\ell} \supset 3$ phases absorbed in l_i
 - $P_{kk}^{\nu} \supset N-1$ phases absorbed in ν_i (only possible if ν_i is Dirac)
- \Rightarrow For $N = 3 + s$: $U_{\text{LEP}} \supset 3(1+s)$ angles + $(2s+1)$ Dirac phases + $(s+2)$ Maj phases

Lepton Mixing

$$U_{\text{LEP}} \equiv 3 \times N \text{ matrix} \quad U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^{\ell} V_L^{\ell \dagger ik} V^{\nu kj} P_{jj}^{\nu}$$

- $P_{ii}^{\ell} \supset 3$ phases absorbed in l_i
 - $P_{kk}^{\nu} \supset N-1$ phases absorbed in ν_i (only possible if ν_i is Dirac)
- \Rightarrow For $N = 3 + s$: $U_{\text{LEP}} \supset 3(1+s)$ angles + $(2s+1)$ Dirac phases + $(s+2)$ Maj phases
- For example for 3 Dirac ν : 3 Mixing angles + 1 Dirac Phase

$$U_{\text{LEP}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{21} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Lepton Mixing

$$U_{\text{LEP}} \equiv 3 \times N \text{ matrix} \quad U_{\text{LEP}}^{ij} = \sum_{k=1}^3 P_{ii}^{\ell} V_L^{\ell \dagger ik} V^{\nu kj} P_{jj}^{\nu}$$

- $P_{ii}^{\ell} \supset 3$ phases absorbed in l_i
 - $P_{kk}^{\nu} \supset N-1$ phases absorbed in ν_i (only possible if ν_i is Dirac)
- \Rightarrow For $N = 3 + s$: $U_{\text{LEP}} \supset 3(1+s)$ angles + $(2s+1)$ Dirac phases + $(s+2)$ Maj phases
- For example for 3 Dirac ν : 3 Mixing angles + 1 Dirac Phase

$$U_{\text{LEP}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{21} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- For 3 Majorana ν : 3 Mixing angles + 1 Dirac Phase + 2 Majorana Phases


$$U_{\text{LEP}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{21} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\phi_2} & 0 \\ 0 & 0 & e^{i\phi_3} \end{pmatrix}$$

Effects of ν Mass: Flavour Transitions

- Flavour (\equiv Interaction) basis (production and detection): ν_e , ν_μ and ν_τ
- Mass basis (free propagation in space-time): ν_1 , ν_2 and $\nu_3 \dots$


Effects of ν Mass: Flavour Transitions

- Flavour (\equiv Interaction) basis (production and detection): ν_e, ν_μ and ν_τ
- Mass basis (free propagation in space-time): ν_1, ν_2 and $\nu_3 \dots$
- In general interaction eigenstates \neq propagation eigenstates



Effects of ν Mass: Flavour Transitions

- Flavour (\equiv Interaction) basis (production and detection): ν_e, ν_μ and ν_τ
- Mass basis (free propagation in space-time): ν_1, ν_2 and $\nu_3 \dots$
- In general interaction eigenstates \neq propagation eigenstates



- The probability $P_{\alpha\beta}$ of producing neutrino with flavour α and detecting with flavour β has to depend on:
 - **Misalignment** between interaction and propagation states ($\equiv U$)
 - **Difference** between propagation **eigenvalues**
 - **Propagation distance**

Mass Induced Flavour Oscillations in Vacuum

- If neutrinos have mass, a weak eigenstate $|\nu_\alpha\rangle$ produced in $l_\alpha + N \rightarrow \nu_\alpha + N'$ is a linear combination of the mass eigenstates ($|\nu_i\rangle$)

$$|\nu_\alpha\rangle = \sum_{i=1}^n U_{\alpha i}^* |\nu_i\rangle$$

U is the leptonic mixing matrix.

Mass Induced Flavour Oscillations in Vacuum

- If neutrinos have mass, a weak eigenstate $|\nu_\alpha\rangle$ produced in $l_\alpha + N \rightarrow \nu_\alpha + N'$ is a linear combination of the mass eigenstates ($|\nu_i\rangle$)

$$|\nu_\alpha\rangle = \sum_{i=1}^n U_{\alpha i}^* |\nu_i\rangle$$

U is the leptonic mixing matrix.

- After a distance L (or time t) it evolves

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^n U_{\alpha i}^* |\nu_i(t)\rangle$$

Mass Induced Flavour Oscillations in Vacuum

- If neutrinos have mass, a weak eigenstate $|\nu_\alpha\rangle$ produced in $l_\alpha + N \rightarrow \nu_\alpha + N'$ is a linear combination of the mass eigenstates ($|\nu_i\rangle$)

$$|\nu_\alpha\rangle = \sum_{i=1}^n U_{\alpha i}^* |\nu_i\rangle$$

U is the leptonic mixing matrix.

- After a distance L (or time t) it evolves

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^n U_{\alpha i}^* |\nu_i(t)\rangle$$

- it can be detected with flavour β with probability

$$P_{\alpha\beta} = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{j=1}^n \sum_{i=1}^n U_{\alpha i}^* U_{\beta j} \langle \nu_j | \nu_i(t) \rangle \right|^2$$

Mass Induced Flavour Oscillations in Vacuum

- The probability

$$P_{\alpha\beta} = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{j=1}^n \sum_{i=1}^n U_{\alpha i}^* U_{\beta j} \langle \nu_j | \nu_i(t) \rangle \right|^2$$

- We call E_i the neutrino energy and m_i the neutrino mass
- Under the approximations:

(1) $|\nu\rangle$ is a *plane wave* $\Rightarrow |\nu_i(t)\rangle = e^{-i E_i t} |\nu_i(0)\rangle$ and using $\langle \nu_j | \nu_i \rangle = \delta_{ij}$

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

with $\Delta_{ij} = (E_i - E_j)t$

Mass Induced Flavour Oscillations in Vacuum

- The probability

$$P_{\alpha\beta} = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{j=1}^n \sum_{i=1}^n U_{\alpha i}^* U_{\beta j} \langle \nu_j | \nu_i(t) \rangle \right|^2$$

- We call E_i the neutrino energy and m_i the neutrino mass
- Under the approximations:

(1) $|\nu\rangle$ is a *plane wave* $\Rightarrow |\nu_i(t)\rangle = e^{-i E_i t} |\nu_i(0)\rangle$ and using $\langle \nu_j | \nu_i \rangle = \delta_{ij}$

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

with $\Delta_{ij} = (E_i - E_j)t$

(2) *relativistic* ν

$$E_i = \sqrt{p_i^2 + m_i^2} \simeq p_i + \frac{m_i^2}{2E_i}$$

Mass Induced Flavour Oscillations in Vacuum

- The probability

$$P_{\alpha\beta} = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{j=1}^n \sum_{i=1}^n U_{\alpha i}^* U_{\beta j} \langle \nu_j | \nu_i(t) \rangle \right|^2$$

- We call E_i the neutrino energy and m_i the neutrino mass
- Under the approximations:

(1) $|\nu\rangle$ is a *plane wave* $\Rightarrow |\nu_i(t)\rangle = e^{-i E_i t} |\nu_i(0)\rangle$ and using $\langle \nu_j | \nu_i \rangle = \delta_{ij}$

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

with $\Delta_{ij} = (E_i - E_j)t$

(2) *relativistic* ν

$$E_i = \sqrt{p_i^2 + m_i^2} \simeq p_i + \frac{m_i^2}{2E_i}$$

(3) Lowest order in mass $p_i \simeq p_j = p \simeq E$

$$\frac{\Delta_{ij}}{2} = \frac{(m_i^2 - m_j^2) L}{4 E} = 1.27 \frac{m_i^2 - m_j^2}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \operatorname{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \operatorname{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

- The first term $\delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right)$ equal for $\bar{\nu}$ ($U \rightarrow U^*$)
 \rightarrow conserves **CP**

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

- The first term $\delta_{\alpha\beta} - 4 \sum_{i < j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right)$ equal for $\bar{\nu}$ ($U \rightarrow U^*$)
→ conserves **CP**
- The last piece $2 \sum_{i < j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$ opposite sign for $\bar{\nu}$
→ violates **CP**

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

- The first term $\delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right)$ equal for $\bar{\nu}$ ($U \rightarrow U^*$)
→ conserves **CP**

- The last piece $2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$ opposite sign for $\bar{\nu}$
→ violates **CP**

- If $\alpha = \beta \Rightarrow \text{Im}[U_{\alpha i} U_{\alpha i}^* U_{\alpha j}^* U_{\alpha j}] = \text{Im}[|U_{\alpha i}^*|^2 |U_{\alpha j}|^2] = 0$

\Rightarrow CP violation observable only for $\beta \neq \alpha$

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

- The first term $\delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right)$ equal for $\bar{\nu}$ ($U \rightarrow U^*$)
→ conserves **CP**
- The last piece $2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$ opposite sign for $\bar{\nu}$
→ violates **CP**

- $P_{\alpha\beta}$ depends on Neutrino Parameters

- $\Delta m_{ij}^2 = m_i^2 - m_j^2$ The mass differences
- $U_{\alpha j}$ The mixing angles (and Dirac phases)

- and on Two set-up Parameters:

- E The neutrino energy
- L Distance ν source to detector

Mass Induced Flavour Oscillations in Vacuum

- The oscillation probability:

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right) + 2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$$

$$\frac{\Delta_{ij}}{2} = \frac{(E_i - E_j)L}{2} = 1.27 \frac{(m_i^2 - m_j^2)}{\text{eV}^2} \frac{L/E}{\text{Km/GeV}}$$

- The first term $\delta_{\alpha\beta} - 4 \sum_{i<j}^n \text{Re}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin^2 \left(\frac{\Delta_{ij}}{2} \right)$ equal for $\bar{\nu}$ ($U \rightarrow U^*$)
→ conserves **CP**
- The last piece $2 \sum_{i<j} \text{Im}[U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}] \sin(\Delta_{ij})$ opposite sign for $\bar{\nu}$
→ violates **CP**

- $P_{\alpha\beta}$ depends on Neutrino Parameters

and on Two set-up Parameters:


- $\Delta m_{ij}^2 = m_i^2 - m_j^2$ The mass differences
- $U_{\alpha j}$ The mixing angles (and Dirac phases)

- E The neutrino energy
- L Distance ν source to detector

- No information on mass scale nor Majorana phases

Dirac or Majorana? ν -less Double- β Decay


Lectures of R. Schwingenheuer



- Amplitude includes
$$[\bar{e}\gamma^\mu L\nu_e][\bar{e}\gamma^\mu L\nu_e] = \sum_{ij} U_{ei}U_{ej}^p [\bar{e}\gamma^\mu \nu_i][\bar{e}\gamma^\mu \nu_j]$$

Dirac or Majorana? ν -less Double- β Decay


Lectures of R. Schwingenheuer



- Amplitude includes $[\bar{e}\gamma^\mu L\nu_e][\bar{e}\gamma^\mu L\nu_e] = \sum_{ij} U_{ei}U_{ej}^p [\bar{e}\gamma^\mu \nu_i][\bar{e}\gamma^\mu \nu_j]$
 - If ν_i Dirac $\Rightarrow \nu_i$ annihilates a neutrino and creates an antineutrino
 \Rightarrow no same state \Rightarrow Amplitude = 0
 - If ν_i Majorana $\Rightarrow \nu_i = \nu_i^c$ annihilates and creates a neutrino=antineutrino
 \Rightarrow same state \Rightarrow Amplitude $\propto \overline{\nu_i}(\nu_i)^T \neq 0$

Dirac or Majorana? ν -less Double- β Decay

Lectures of R. Schwingenheuer



- Amplitude includes $[\bar{e}\gamma^\mu L\nu_e][\bar{e}\gamma^\mu L\nu_e] = \sum_{ij} U_{ei}U_{ej}^p [\bar{e}\gamma^\mu \nu_i][\bar{e}\gamma^\mu \nu_j]$
 - If ν_i Dirac $\Rightarrow \nu_i$ annihilates a neutrino and creates an antineutrino
 \Rightarrow no same state \Rightarrow Amplitude = 0
 - If ν_i Majorana $\Rightarrow \nu_i = \nu_i^c$ annihilates and creates a neutrino=antineutrino
 \Rightarrow same state \Rightarrow Amplitude $\propto \nu_i (\nu_i)^T \neq 0$

- If Majorana m_ν only source of L -violation

\Rightarrow Amplitude of ν -less- $\beta\beta$ decay is proportional to $\langle m_{ee} \rangle = \sum_j U_{ej}^2 m_j$

Neutrino Mass Scale: Tritium β Decay

- Fermi proposed a kinematic search of ν_e mass from beta spectra in ${}^3\text{H}$ beta decay



- For “allowed” nuclear transitions, the electron spectrum is given by phase space alone

$$K(T) \equiv \sqrt{\frac{dN}{dT} \frac{1}{C_p E F(E)}} \propto \sqrt{(Q - T) \sqrt{(Q - T)^2 - m_{\nu_e}^2}}$$

$T = E_e - m_e$, Q = maximum kinetic energy, (for ${}^3\text{H}$ beta decay $Q = 18.6$ KeV)

Taking into account mixing

$$m_{\nu_e}^{\text{eff}} \equiv \sqrt{\sum m_{\nu_j}^2 |U_{ej}|^2}$$

Neutrino Mass Scale: Tritium β Decay

- Fermi proposed a kinematic search of ν_e mass from beta spectra in ${}^3\text{H}$ beta decay



- For “allowed” nuclear transitions, the electron spectrum is given by phase space alone

$$K(T) \equiv \sqrt{\frac{dN}{dT} \frac{1}{C_p E F(E)}} \propto \sqrt{(Q - T) \sqrt{(Q - T)^2 - m_{\nu_e}^2}}$$

$T = E_e - m_e$, Q = maximum kinetic energy, (for ${}^3\text{H}$ beta decay $Q = 18.6$ KeV)


Taking into account mixing

$$m_{\nu_e}^{\text{eff}} \equiv \sqrt{\sum m_{\nu_j}^2 |U_{ej}|^2}$$

- $m_{\nu} \neq 0 \Rightarrow$ distortion from the straight-line at the end point of the spectrum

$$m_{\nu} = 0 \Rightarrow T_{\text{max}} = Q$$

$$m_{\nu} \neq 0 \Rightarrow T_{\text{max}} = Q - m_{\nu}$$



Neutrino Mass Scale: Tritium β Decay

- Fermi proposed a kinematic search of ν_e mass from beta spectra in ${}^3\text{H}$ beta decay



- For “allowed” nuclear transitions, the electron spectrum is given by phase space alone

$$K(T) \equiv \sqrt{\frac{dN}{dT} \frac{1}{C_p E F(E)}} \propto \sqrt{(Q - T) \sqrt{(Q - T)^2 - m_{\nu_e}^2}}$$

$T = E_e - m_e$, Q = maximum kinetic energy, (for ${}^3\text{H}$ beta decay $Q = 18.6$ KeV)


Taking into account mixing

$$m_{\nu_e}^{\text{eff}} \equiv \sqrt{\sum m_{\nu_j}^2 |U_{ej}|^2}$$

- $m_{\nu} \neq 0 \Rightarrow$ distortion from the straight-line at the end point of the spectrum

$$m_{\nu} = 0 \Rightarrow T_{\text{max}} = Q$$

$$m_{\nu} \neq 0 \Rightarrow T_{\text{max}} = Q - m_{\nu}$$




– At present only a bound: $m_{\nu_e}^{\text{eff}} < 0.8$ eV (at 90 % CL) (Katrín)

– Katrín operating can improve present sensitivity to $m_{\nu_e}^{\text{eff}} \sim 0.3$ eV

Probes of Mass Scale

Single β decay : Pure kinematics, **Dirac** or **Majorana** ν 's, only model independent




$$m_{\nu_e}^2 = \sum m_j^2 |U_{ej}|^2$$

Present bound: $m_{\nu_e} \leq 0.8$ eV (90% CL KATRIN 2021)

Katrin (20XX) Sensitivity to $m_{\nu_e} \sim 0.2$ eV

ν -less Double- β decay: \Leftrightarrow **Majorana ν 's**



If m_{ν} only source of ΔL $T_{1/2}^{0\nu} = \frac{m_e}{G_{0\nu} M_{\text{nucl}}^2 m_{ee}^2}$

$$m_{ee} = \left| \sum U_{ej}^2 m_j \right|$$

Present Bounds: $m_{ee} < 0.04 - 0.2$ eV

COSMO for **Dirac** or **Majorana**

$$\sum m_i$$

Lecture S. Pastor

Back up Slides

The Other Flavours

ν coming out of a nuclear reactor is $\bar{\nu}_e$ because it is emitted together with an e^-

Question: Is it different from the muon type neutrino ν_μ that could be associated to the muon? Or is this difference a theoretical arbitrary convention?


The Other Flavours

ν coming out of a nuclear reactor is $\bar{\nu}_e$ because it is emitted together with an e^-

Question: Is it different from the muon type neutrino ν_μ that could be associated to the **muon**? Or is this difference a theoretical arbitrary convention?

In 1959 **M. Schwartz** thought of producing an intense ν beam from π 's decay (produced when a proton beam of GeV energy hits matter)

Schwartz, Lederman, Steinberger and **Gaillard** built a spark chamber (a 10 tons of neon gas) to detect ν_μ




The Other Flavours

ν coming out of a nuclear reactor is $\bar{\nu}_e$ because it is emitted together with an e^-

Question: Is it different from the muon type neutrino ν_μ that could be associated to the **muon**? Or is this difference a theoretical arbitrary convention?

In 1959 **M. Schwartz** thought of producing an intense ν beam from π 's decay (produced when a proton beam of GeV energy hits matter)

Schwartz, Lederman, Steinberger and **Gaillard** built a spark chamber (a 10 tons of neon gas) to detect ν_μ



They observe 40 ν interactions: in 6 an e^- comes out and in 34 a μ^- comes out.

If $\nu_\mu \equiv \nu_e \Rightarrow$ equal numbers of μ^- and e^-


The Other Flavours

ν coming out of a nuclear reactor is $\bar{\nu}_e$ because it is emitted together with an e^-

Question: Is it different from the muon type neutrino ν_μ that could be associated to the **muon**? Or is this difference a theoretical arbitrary convention?

In 1959 **M. Schwartz** thought of producing an intense ν beam from π 's decay (produced when a proton beam of GeV energy hits matter)

Schwartz, Lederman, Steinberger and **Gaillard** built a spark chamber (a 10 tons of neon gas) to detect ν_μ



They observe 40 ν interactions: in 6 an e^- comes out and in 34 a μ^- comes out.

If $\nu_\mu \equiv \nu_e \Rightarrow$ equal numbers of μ^- and $e^- \Rightarrow$ **Conclusion: ν_μ is a different particle**


The Other Flavours

ν coming out of a nuclear reactor is $\bar{\nu}_e$ because it is emitted together with an e^-

Question: Is it different from the muon type neutrino ν_μ that could be associated to the **muon**? Or is this difference a theoretical arbitrary convention?

In 1959 **M. Schwartz** thought of producing an intense ν beam from π 's decay (produced when a proton beam of GeV energy hits matter)

Schwartz, Lederman, Steinberger and **Gaillard** built a spark chamber (a 10 tons of neon gas) to detect ν_μ



They observe 40 ν interactions: in 6 an e^- comes out and in 34 a μ^- comes out.

If $\nu_\mu \equiv \nu_e \Rightarrow$ equal numbers of μ^- and $e^- \Rightarrow$ **Conclusion: ν_μ is a different particle**

In 1977 **Martin Perl** discovers the particle tau \equiv the third lepton family.


The ν_τ was observed by **DONUT** experiment at FNAL in 1998 (officially in Dec. 2000).

Neutrino Helicity

Neutrino Helicity

- The neutrino helicity was measured in 1957 in a experiment by Goldhaber et al.

- Using the electron capture reaction




with $J({}^{152}\text{Eu}) = J({}^{152}\text{Sm}) = 0$ and $L(e^{-}) = 0$

Neutrino Helicity

- The neutrino helicity was measured in 1957 in a experiment by Goldhaber et al.

- Using the electron capture reaction



with $J({}^{152}\text{Eu}) = J({}^{152}\text{Sm}) = 0$ and $L(e^{-}) = 0$


- Angular momentum conservation \Rightarrow

$$\left\{ \begin{array}{lcl} J_z(e^{-}) & = & J_z(\nu) + J_z(\text{Sm}^{*}) \\ & = & J_z(\nu) + J_z(\gamma) \\ \pm \frac{1}{2} & = & \mp \frac{1}{2} \quad \pm 1 \Rightarrow J_z(\nu) = -\frac{1}{2} J_z(\gamma) \end{array} \right.$$

Neutrino Helicity

- The neutrino helicity was measured in 1957 in a experiment by Goldhaber et al.

- Using the electron capture reaction



with $J({}^{152}\text{Eu}) = J({}^{152}\text{Sm}) = 0$ and $L(e^{-}) = 0$

- Angular momentum conservation \Rightarrow

$$\begin{cases} J_z(e^{-}) &= J_z(\nu) + J_z(\text{Sm}^{*}) \\ &= J_z(\nu) + J_z(\gamma) \\ \pm \frac{1}{2} &= \mp \frac{1}{2} \quad \pm 1 \Rightarrow J_z(\nu) = -\frac{1}{2} J_z(\gamma) \end{cases}$$


- Nuclei are heavy $\Rightarrow \vec{p}({}^{152}\text{Eu}) \simeq \vec{p}({}^{152}\text{Sm}) \simeq \vec{p}({}^{152}\text{Sm}^{*}) = 0$

So momentum conservation $\Rightarrow \vec{p}(\nu) = -\vec{p}(\gamma) \Rightarrow \nu \text{ helicity} = \gamma \text{ helicity}$

Neutrino Helicity

- The neutrino helicity was measured in 1957 in a experiment by Goldhaber et al.

- Using the electron capture reaction



with $J({}^{152}\text{Eu}) = J({}^{152}\text{Sm}) = 0$ and $L(e^{-}) = 0$

- Angular momentum conservation \Rightarrow

$$\begin{cases} J_z(e^{-}) &= J_z(\nu) + J_z(\text{Sm}^{*}) \\ &= J_z(\nu) + J_z(\gamma) \\ \pm \frac{1}{2} &= \mp \frac{1}{2} \quad \pm 1 \Rightarrow J_z(\nu) = -\frac{1}{2} J_z(\gamma) \end{cases}$$


- Nuclei are heavy $\Rightarrow \vec{p}({}^{152}\text{Eu}) \simeq \vec{p}({}^{152}\text{Sm}) \simeq \vec{p}({}^{152}\text{Sm}^{*}) = 0$

So momentum conservation $\Rightarrow \vec{p}(\nu) = -\vec{p}(\gamma) \Rightarrow \nu \text{ helicity} = \gamma \text{ helicity}$

- Goldhaber et al found γ had negative helicity $\Rightarrow \nu$ has helicity -1

Number of Neutrinos


- The counting of **light left-handed neutrinos** is based on the family structure of the SM assuming a universal diagonal NC coupling:



$$j_Z^\mu = \sum_{\alpha} \bar{\nu}_{\alpha L} \gamma^\mu \nu_{\alpha L}$$

Number of Neutrinos

- The counting of **light left-handed neutrinos** is based on the family structure of the SM assuming a universal diagonal NC coupling:



$$j_Z^\mu = \sum_{\alpha} \bar{\nu}_{\alpha L} \gamma^\mu \nu_{\alpha L}$$

- For $m_{\nu_i} < m_Z/2$ one can use the total Z -width Γ_Z to extract N_ν


$$N_\nu = \frac{\Gamma_{\text{inv}}}{\Gamma_\nu} \equiv \frac{1}{\Gamma_\nu} (\Gamma_Z - \Gamma_h - 3\Gamma_\ell)$$

$$= \frac{\Gamma_\ell}{\Gamma_\nu} \left[\sqrt{\frac{12\pi R_{h\ell}}{\sigma_h^0 m_Z^2}} - R_{h\ell} - 3 \right]$$

Γ_{inv} = the invisible width

Γ_h = the total hadronic width

Γ_ℓ = width to charged lepton




Leads $N_\nu = 2.9840 \pm 0.0082$

Neutrinos always “Left-Handed” \equiv Massless

- If ν had a mass they would not go to the speed of light:
 \Rightarrow the direction of its momentum depends on the reference frame


So in one reference frame




Neutrinos always “Left-Handed” \equiv Massless

- If ν had a mass they would not go to the speed of light:
 \Rightarrow the direction of its momentum depends on the reference frame

So in one reference frame




And in another



Neutrinos always “Left-Handed” \equiv Massless

- If ν had a mass they would not go to the speed of light:
 \Rightarrow the direction of its momentum depends on the reference frame


So in one reference frame



So if always left \Rightarrow

Strictly massless

And in another



ν Mass from Non-Renormalizable Operator

ν Mass from Non-Renormalizable Operator

If SM is an effective low energy theory, for $E \ll \Lambda_{\text{NP}}$

- The same particle content as the SM and same pattern of symmetry breaking
- But there can be non-renormalizable
(dim > 4) operators

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_n \frac{1}{\Lambda_{\text{NP}}^{n-4}} \mathcal{O}_n$$

ν Mass from Non-Renormalizable Operator

If SM is an effective low energy theory, for $E \ll \Lambda_{\text{NP}}$

- The same particle content as the SM and same pattern of symmetry breaking
- But there can be non-renormalizable (dim > 4) operators

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_n \frac{1}{\Lambda_{\text{NP}}^{n-4}} \mathcal{O}_n$$

First NP effect \Rightarrow dim=5 operator

There is only one!

$$\mathcal{L}_5 = \frac{Z_{ij}^\nu}{\Lambda_{\text{NP}}} \left(\overline{L_{L,i}} \tilde{\phi} \right) \left(\tilde{\phi}^T L_{L,j}^C \right)$$

ν Mass from Non-Renormalizable Operator

If SM is an effective low energy theory, for $E \ll \Lambda_{\text{NP}}$

- The same particle content as the SM and same pattern of symmetry breaking
- But there can be non-renormalizable
(dim > 4) operators

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_n \frac{1}{\Lambda_{\text{NP}}^{n-4}} \mathcal{O}_n$$

First NP effect \Rightarrow dim=5 operator

There is only one!

$$\mathcal{L}_5 = \frac{Z_{ij}^\nu}{\Lambda_{\text{NP}}} \left(\overline{L_{L,i}} \tilde{\phi} \right) \left(\tilde{\phi}^T L_{L,j}^C \right)$$

which after symmetry breaking

induces a ν Majorana mass

$$(M_\nu)_{ij} = Z_{ij}^\nu \frac{v^2}{\Lambda_{\text{NP}}}$$

ν Mass from Non-Renormalizable Operator

If SM is an effective low energy theory, for $E \ll \Lambda_{\text{NP}}$

- The same particle content as the SM and same pattern of symmetry breaking
- But there can be non-renormalizable
(dim > 4) operators

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_n \frac{1}{\Lambda_{\text{NP}}^{n-4}} \mathcal{O}_n$$

First NP effect \Rightarrow dim=5 operator

There is only one!

$$\mathcal{L}_5 = \frac{Z_{ij}^\nu}{\Lambda_{\text{NP}}} \left(\overline{L_{L,i}} \tilde{\phi} \right) \left(\tilde{\phi}^T L_{L,j}^C \right)$$

which after symmetry breaking

induces a ν Majorana mass

$$(M_\nu)_{ij} = Z_{ij}^\nu \frac{v^2}{\Lambda_{\text{NP}}}$$

Implications:

- It is natural that ν mass is the first evidence of NP
- Naturally $m_\nu \ll$ other fermions masses $\sim \lambda^f v$ if $\Lambda_{\text{NP}} \gg v$
- See-saw with heavy ν_R integrated out is a particular example of this