

# Study of semitauonic B-meson decays at Belle and Belle II

---

Mateusz Kaleta

Institute of Nuclear Physics Polish Academy of Sciences, Krakow


June 2024

- Introduction and motivation
- Overview of current measurements
- Belle (II) Experiment
- Analysis
- Summary

## Motivation


- $B \rightarrow D^{(*)} \tau \nu$  decays are sensitive to new amplitudes at tree-level, heavy lepton in the final state
- Large number of observables:  $R(D^{(*)})$ , polarisations  $\tau$  in  $D^*$ ,  $q^2$  distributions,
- Good theoretical tools; precise SM predictions, small hadronic uncertainties.

$$R(D^{(*)}) = \frac{\mathcal{B}(B \rightarrow D^{(*)} \tau \nu)}{\mathcal{B}(B \rightarrow D^{(*)} \ell \nu)}$$




Combined  $R(D)$  and  $R(D^*)$  in tension with SM prediction at  $3\sigma$  level.

# Introduction



[M. Tanaka, R.Watanabe, New physics in the weak interaction of  $B \rightarrow D^{(*)} \tau \nu$ ]

## $D^*$ polarisation at Belle



- The signal yields obtained in the bins of  $\cos\theta_{hel}$  were re-weighted with the following scale factors ( $s_i$ ) to correct for acceptance variations.
- Correction factors  $s_i$  extracted from MC **assuming Standard Model decay dynamics**

$\cos\theta_{hel}$	$s$
$(-1, -0.67)$	$0.98 \pm 0.01$
$(-0.67, -0.33)$	$0.96 \pm 0.01$
$(-0.33, 0)$	$1.08 \pm 0.01$

$$B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$$

$$\tau^+ \rightarrow \ell^+ \nu_\ell \bar{\nu}_\tau, \tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$$

$$F_L(D^*) = 0.60 \pm 0.08 \text{ (stat)} \pm 0.04 \text{ (syst)}$$

Results consistent with SM prediction at  $1.6\sigma - 1.8\sigma$

Karol Adamczyk. PhD thesis, [arXiv:1903.03102] (Belle Collaboration)

## $D^*$ polarisation at LHCb (2023)

$$B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$$

$$\tau^+ \rightarrow \pi^+ \pi^- \pi^+ (\pi^0) \nu_\tau$$

$$F_L(D^*) = 0.43 \pm 0.06 \text{ (stat)} \pm 0.03 \text{ (syst)}$$

Compatible with SM predictions and with Belle results.

[arXiv:2311.05224v1] (LHCb Collaboration)

## $\tau$ polarisation at Belle

$$B \rightarrow \bar{D}^* \tau^+ \nu_\tau$$

$$\tau^- \rightarrow \pi^- \nu_\tau, \rho^- \nu_\tau$$

$$P_\tau = -0.38 \pm 0.51 \text{ (stat)} \pm 0.20 \text{ (syst)}$$

Consistent with SM prediction at  $0.6\sigma$   
[PRL118 211801 (2017), PRD97 012004 (2018)]  
(Belle Collaboration)

Both measurements performed assuming Standard Model decay dynamics.

# Goal of this analysis


## Main goal

- Enhancing experimental constraints on  $B \rightarrow \bar{D}^* \tau \nu \tau$  by precise measurements of angular observables.
- Focusing on  $F_L(D^*)$ .

## Specific goals

- **Model-independent corrections for acceptance effects**
- Increase statistics w.r.t. previous Belle analysis:
  - combined analysis of Belle and Belle II data
  - adding charged B channel:  $B^+ \rightarrow D^* \tau \nu$
  - including more D decay channels in the analysis
- Perform measurements in several  $q^2$  bins

# Kinematic variables



- $q^2$  - effective mass squared of the  $\tau\nu$  system
- $\theta_{hel}(D^*)$  - angle between D and B in  $D^*$  rest frame



$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_{hel}(D^*)} = \frac{3}{4} \left[ 2F_L(D^*) \cos^2\theta_{hel}(D^*) + (1 - F_L(D^*)) \sin^2\theta_{hel}(D^*) \right]$$

$q^2$  and  $\cos\theta_{hel}(D^*)$  can be reconstructed at B-factories with hadronic decays of  $B_{tag}$





# Acceptance effects

Belle



Belle II



Generated (black) vs. reconstructed\* (red)  $q^2$  and  $\cos\theta_{hel}(D^*)$  distributions for Belle (top) and Belle II (bottom).

\*True kinematics is used for reconstructed events.

Decay channel:  $\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau$   
 $D^{*+} \rightarrow D^0 \pi^+$   
 $\tau^- \rightarrow \ell^- \bar{\nu}_\ell \nu_\tau$

Signal decays generated assuming Standard Model decay dynamics.

# Efficiency map

1. Pick four variables that characterize the decay and can be reconstructed experimentally:


- $\cos\theta_{hel}(D^*)$  - cosine helicity angle  $D^*$
- $q^2$  - four-momentum transfer squared
- $\tilde{E}_d$  - normalised  $\tau$  daughter energy
- $\cos\theta_d$  -  $\tau$  daughter polar angle

2. Create a 4D efficiency map by dividing reconstructed histograms by generated ones.

$$W_{ijkl} = \frac{N_{ijkl}^{rec}}{N_{ijkl}^{gen}} \frac{N_{total}^{gen}}{N_{total}^{rec}}$$


$N_{ijkl}$  - number of events per bin

$N_{tot}$  - total number of events



1D projections of 4D efficiency maps. Each variable was divided in 5 equidistant bins. Plot generated for Belle geometry, using Standard Model decay dynamics.

3. Reweight reconstructed distributions using  $w_{ijkl}$  to recover generated observables.



Generated (red) and reconstructed + reweighted distributions (blue). Plots made on independent sample generated with non-SM decay dynamics.

- Semitauonic B-meson decays currently on spotlight
- Improving experimental results for angular analyses can be useful for interpretation of current anomalies
- Studies on signal MC show the  $D^*$  polarisation measurement is challenging due to large acceptance effects
- We plan to apply model-independent acceptance corrections not considered previously